



Quick Example of Benefits of Pairing Trajectory Accuracy Data

By

Mike Paglione

July 24, 2000





What is the Goal?

Determine that if running in interfacility mode with ZME and ZID, URET DU has a statistically different horizontal trajectory error than running with ZME in single site mode (with at least 0.05 confidence).





Description of Experiment

- Used 1 hour Initial Delivery Scenario
 - Ran ZME/ZID interfacility run
 - Ran ZME single site run
- Collected each of ZME's SSGs and Ran Trajectory Accuracy Analysis
 - Used all error horizontal accuracy data
 - Facilitated by Oracle 'select' command¹


¹ select avg(horz_err) from traj_metrics group by look_ahead_tm, acid_cid.





Compare Distributions Graphically

IFA VS. SINGLE HORZ ERROR (At Look Ahead Zero Only)





Compare Distributions Statistically

- Sample Mean per Flight at Look Ahead Time of Zero
- Sample Statistics for IFA
 - Mean Horizontal Error = 1.35 nm
 - Standard Deviation of Horz. Error = 0.92 nm
 - Sample Quantity, $n = 220$ flights
- Sample Statistics for Single Site
 - Mean Horizontal Error = 1.23 nm
 - Standard Deviation of Horz. Error = 0.60 nm
 - Sample Quantity, $n = 220$ flights





Perform Hypothesis Test

- On means assuming unequal variance
- Use Smith-Satterthwaite Test (Devore, 1987)
 - Test Statistic = $t' = [\text{Avg}(\text{IFA}) - \text{Avg}(\text{Single})] / [\text{sqrt}(\text{Stddev}(\text{IFA})^2/n + \text{Stddev}(\text{Single})^2/n)]$
 - Reject Hypothesis if $t' \geq t_{0.025, \text{dof}}$ or $t' \leq -t_{0.025, \text{dof}}$
 - DOF = degrees of freedom ~
- Results
 - $t' = 1.615$
 - $t_{0.025, 379} = 1.966$
 - P-value = 0.107
- Conclusion: No Significant Difference between Means at 0.05 Confidence





Alternative Hypothesis Test

- Test Difference between Flights of IFA vs Single
- Apply Paired Data Test
 - Calculate difference , d, for each flight/look ahead
 - Sample Quantity of d's = n
 - Test Statistic = $t_p = [\text{avg}(d)] / [\text{stddev}(d) / \text{sqrt}(n)]$
 - Reject Hypothesis if $t_p \geq t_{0.025, n-1}$ or $t_p \leq -t_{0.025, n-1}$
 - DOF = degrees of freedom = $n - 1$
- Results
 - $t_p = 2.29$ and $n = 220$
 - $t_{0.025, 221} = 1.97$
 - P-value = 0.023
- Conclusion: Is a Significant Difference with 0.05 Confidence, Reject Hypothesis!





Why Difference in Results?

- “If there is a great heterogeneity between subjects (large variance) and a large correlation within subjects, then the loss in degrees of freedom will be compensated for by the increased precision associated with pairing ...” (Devore, 1987)
- Flight Accuracy Between Runs not Independent
- Variance is Large Between Flights





Pairing Improves Precision

➤ Compare Each Confidence Interval

- Paired CI

- ✓ $\text{Avg}(d) \pm t_{0.025, n-1} (\text{stddev}(d) / \sqrt{n})$

- ✓ $0.012 \pm 0.103 = [0.02, 0.22]$

- Pooled or Independent Analysis CI

- ✓ $[\text{Avg}(\text{ifa}) - \text{Avg}(\text{single})] \pm t_{0.025, v} [\sqrt{\text{Stddev}(\text{IFA})^2/n + \text{Stddev}(\text{Single})^2/n}]$

- ✓ $0.012 \pm 0.146 = [-.03, 0.27]$

➤ Conclusions

- Any decision made on independent CI is a third less precise!
- Independent CI includes error between flights as well as between runs, while paired CI blocks between flight error and focuses on runs only!





References

- Devore, Jay L., *Probability and Statistics for Engineering and the Sciences*, Second Edition, 1987.
- Montgomery, Douglas C., *Design and Analysis of Experiments*, Fourth Edition, 1997.

